



HAL
open science

A two-particle, four-mode interferometer for atoms

Pierre I Dussarrat, Maxime I Perrier, Almazbek I Imanaliev, Raphael Lopes,
Alain Aspect, Marc Cheneau, Denis Boiron, Christoph I Westbrook

► **To cite this version:**

Pierre I Dussarrat, Maxime I Perrier, Almazbek I Imanaliev, Raphael Lopes, Alain Aspect, et al.. A two-particle, four-mode interferometer for atoms. 2017. hal-01551704v1

HAL Id: hal-01551704

<https://hal-iogs.archives-ouvertes.fr/hal-01551704v1>

Preprint submitted on 30 Jun 2017 (v1), last revised 17 Oct 2017 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A two-particle, four-mode interferometer for atoms

Pierre Dussarrat,¹ Maxime Perrier,¹ Almazbek Imanaliev,² Raphael Lopes,³
Alain Aspect,¹ Marc Cheneau,¹ Denis Boiron,¹ and Christoph I. Westbrook¹

¹*Laboratoire Charles Fabry, Institut d'Optique Graduate School,
CNRS, Université ParisSaclay, 91120 Palaiseau, France*

²*LNE-SYRTE, Observatoire de Paris, PSL Research University, CNRS,
Sorbonne Universités, UPMC Université Paris 06, 75014 Paris, France*

³*Cavendish Laboratory, University of Cambridge, Cambridge CB3 0HE, United Kingdom*

We present a free-space interferometer to observe two-particle interference of a pair of atoms with entangled momenta. The source of atom pairs is a Bose–Einstein condensate subject to a dynamical instability, and the interferometer is realized using Bragg diffraction on optical lattices, in the spirit of our recent Hong–Ou–Mandel experiment. We report an observation consistent with an entangled state at the input of the interferometer. We explain how our current setup can be extended to enable a test of a Bell inequality on momentum observables.

A key element of the second quantum revolution [1, 2] is entanglement [3]. Its extraordinary character comes from the fact that the many-body wave-function of entangled particles can only be described in a configuration space associated with the tensor product of the configuration spaces of the individual particles. When one insists on describing it in our ordinary space-time, one has to face the problem of non-locality [4–6]. This is clearly illustrated by the violation of Bell’s inequalities [7], which apply to any system that can be described in the spirit of the local realist worldview of Einstein, in which physical reality lies in our ordinary space-time [8].

While the violation of Bell’s inequalities stems from two-particle interferences observed with entangled pairs, the converse is not true: not all phenomena associated with two-particle interference can lead to a violation of Bell’s inequalities. This is for instance the case of the Hanbury Brown–Twiss effect for thermal bosons [9, 10], or the Hong–Ou–Mandel effect [11]: the quantum description appeals to two-particle interference but no non-locality is involved. This is because the latter effects involve only two modes for two indistinguishable particles, while a configuration leading to the violation of Bell’s inequalities requires four modes that can be made to interfere two by two in different places [12].

Ever more ideal experimental tests of Bell’s inequalities have been performed with low energy photons, internal states of trapped ions and nitrogen-vacancy centers (see references in [13, 14]). But we know of no experiments on two-particle interference in four modes associated with motional degrees of freedom (position or momentum), in a configuration permitting, even in principle, a Bell inequality test [15]. Such tests involving mechanical observables are desirable, in particular because they may allow one to touch upon the interface between quantum mechanics and gravitation [16].

In this paper, we present a setup to measure two-particle interferences with atoms entangled in momentum. We observe correlations consistent with the presence of an entangled state, and discuss how the appara-




FIG. 1. Diagram of a two-particle, four-mode interferometer. An atom pair in the entangled momentum state (1) is emitted at time $t = 0$. Using Bragg diffraction on optical lattices, the four input modes are then deflected at time t_1 , and mixed two by two on the splitters A and B at time $t_2 = 2t_1$. The interference is read-out by detecting the atoms in the output modes A_{\pm} , B_{\pm} , and measuring the probabilities of joint detection $\mathcal{P}(A_{\pm}, B_{\pm})$. The Bragg deflector and splitters differ from their optical analogs, because rather than reversing the incident momentum, they add a fixed momentum $\pm\hbar k_{\ell}$ along the z -axis, where k_{ℓ} is the reciprocal lattice vector. The dashed lines show the Hong–Ou–Mandel configuration.

tus can be extended to realize a test of a Bell inequality. To understand the experiment, consider a Bell state consisting of a pair of atoms in a superposition of distinct momentum modes labeled in the center-of-mass reference frame by $\pm p$ and $\pm p'$:

$$|\Psi\rangle = \frac{1}{\sqrt{2}} (|p, -p\rangle + |p', -p'\rangle) . \quad (1)$$

To probe the coherent superposition of the pair states

$|p, -p\rangle$ and $|p', -p'\rangle$, we use the two-atom interferometer shown in Fig. 1. An analogous interferometer for photons was proposed in Ref. [17], implemented in Ref. [18], and resulted in a Bell inequality violation. Similar configurations for atoms were also analyzed in Refs. [19, 20].

The input modes p and $-p'$ of our interferometer are deflected and mixed on the 50:50 splitter A . Similarly, the input modes p' and $-p$ are deflected and mixed on the 50:50 splitter B . The deflection and mixing are realized with Bragg diffraction on optical lattices. The deflecting lattice is common to the four input modes and is applied at time t_1 . The splitting lattices A and B are applied at time $t_2 = 2t_1$ (the time origin is set at the instant of pair emission). The four output modes of the interferometer, A_{\pm} and B_{\pm} , can be written in terms of the four input modes [21]:

$$|A_+\rangle = \frac{-1}{\sqrt{2}} \left(e^{-i(\phi_A - \phi_D)} |p\rangle + i e^{-i\phi_D} | -p'\rangle \right), \quad (2)$$

$$|A_-\rangle = \frac{-1}{\sqrt{2}} \left(i e^{i\phi_D} |p\rangle + e^{i(\phi_A - \phi_D)} | -p'\rangle \right), \quad (3)$$

$$|B_+\rangle = \frac{-1}{\sqrt{2}} \left(e^{-i(\phi_B - \phi_D)} |p'\rangle + i e^{-i\phi_D} | -p\rangle \right), \quad (4)$$

$$|B_-\rangle = \frac{-1}{\sqrt{2}} \left(i e^{i\phi_D} |p'\rangle + e^{i(\phi_B - \phi_D)} | -p\rangle \right). \quad (5)$$

Here, the phases ϕ_D , ϕ_A and ϕ_B are the relative phases of the laser beams forming the deflecting lattice (ϕ_D) and the splitting lattices (ϕ_A and ϕ_B); they can in principle be separately controlled. We have omitted overall phase factors due to propagation.

Inverting equations (2-5), one readily obtains the expression of the entangled state (1) at the output of the interferometer, which solely depends on ϕ_A and ϕ_B :

$$|\Psi_{\text{out}}\rangle = \frac{1}{2\sqrt{2}} \left[-i(e^{i\phi_A} + e^{i\phi_B})|A_+, B_+\rangle + (e^{i(\phi_A - \phi_B)} - 1)|A_+, B_-\rangle + (e^{-i(\phi_A - \phi_B)} - 1)|A_-, B_+\rangle - i(e^{-i\phi_A} + e^{-i\phi_B})|A_-, B_-\rangle \right]. \quad (6)$$

The probabilities of joint detection in the output modes are given by the squared modulus of the complex amplitudes of the corresponding pair states:

$$\mathcal{P}(A_+, B_+) = \mathcal{P}(A_-, B_-) = \frac{1}{2} \cos^2 [(\phi_A - \phi_B)/2], \quad (7)$$

$$\mathcal{P}(A_+, B_-) = \mathcal{P}(A_-, B_+) = \frac{1}{2} \sin^2 [(\phi_A - \phi_B)/2], \quad (8)$$

while the probabilities of single detection are all equal to 1/2. Observing the modulation of the joint detection probabilities as a function of the phase difference ($\phi_A - \phi_B$) would demonstrate that the initial state is entangled. If rather, we have initially a statistical mixture of the pair states $|p, -p\rangle$ and $|p', -p'\rangle$, there would be no modulation and the probabilities of joint detection would all be equal

to 1/4. The four joint detection probabilities can also be combined in a single correlation coefficient:

$$E = \mathcal{P}(A_+, B_+) + \mathcal{P}(A_-, B_-) \quad (9)$$

$$- \mathcal{P}(A_+, B_-) - \mathcal{P}(A_-, B_+) = V \cos(\phi_A - \phi_B). \quad (10)$$

The visibility V is equal to unity for the input state (1), but it may be reduced in a real experiment due for example to decoherence, or the presence of additional pairs. Of course, a Bell's inequality test remains possible provided $V > 1/\sqrt{2}$ [22].

We now come to our experimental realization. A gaseous Bose-Einstein condensate (BEC) containing 7×10^4 Helium-4 atoms in the metastable $2^3S_1, m_J = 1$ electronic state is confined in an ellipsoidal optical trap with its long axis along the vertical (z) direction. The emission of atom pairs occurs in the presence of a vertical, moving optical lattice formed by the interference of two laser beams with slightly different frequencies [21]. It results from the scattering of two atoms from the BEC and can be thought of as a spontaneous, degenerate four-wave mixing process [23]. The lattice is switched on and off adiabatically in 100 μs , and is maintained at a constant depth for 600 μs . The lattice hold time is tuned to obtain a typical mode occupancy of 0.2 [24]. The probability to emit multiple pairs in the same modes is about 5 times smaller. The optical trap is switched off abruptly as soon as the lattice depth is returned to zero. The atoms are then transferred to the magnetically insensitive $m_J = 0$ state with a two-photon Raman transition and fall freely under the sole influence of gravity. They end their fall on a micro-channel plate detector located 46 cm below the position of the optical trap [25]. The detector records the impact of each atom with an efficiency $\sim 25\%$. We store the arrival times and horizontal positions (x - y -plane), and reconstruct the initial three-dimensional velocity distribution of the atoms.

In Fig. 2, we show the initial velocity distribution of the emitted atom pairs in the y - z -plane. Here, and in the rest of the article, velocities are expressed in the center-of-mass reference frame of the free-falling pairs. The distribution is bimodal, and symmetric under rotation about the z -axis, reflecting the one-dimensional character of the pair emission. We do observe, however, a slight asymmetry in the height of the two maxima. We attribute this asymmetry to momentum-dependent losses occurring during the short time when the emitted atoms spatially overlap with the BEC.

The pairwise emission process is characterized by the normalized cross-correlation:

$$g^{(2)}(v_z^+, v_z^-) = \frac{\langle n(v_z^+) n(v_z^-) \rangle}{\langle n(v_z^+) \rangle \langle n(v_z^-) \rangle}, \quad (11)$$

where $n(v_z^{\pm})$ represents the number of atoms with a velocity $v_z^+ > 0$, or $v_z^- < 0$, along the z -axis and 0 along




FIG. 2. Initial velocity distribution of the emitted atom pairs in the y - z -plane. The color scale represents the total number of atoms detected over 1 169 repetitions of the experiment inside a small integration volume [21]. The velocities are defined with respect to the center-of-mass velocity of the atom pairs, which was measured to be 0, 0 and 94 mm/s along the x , y and z directions, respectively.

the x - and y -axes. Experimentally, we measure this correlation by counting the number of detected atoms inside two small volumes in velocity-space [21], and averaging their product over many realizations (as denoted by $\langle \cdot \rangle$). The correlation obtained in the experiment is displayed in Fig. 3. A two-particle correlation centered around $v_z^+ = -v_z^- \simeq 25$ mm/s is clearly visible and confirms that atoms are indeed emitted in pairs with opposite velocities. Because the pair emission fulfills the quasi-momentum conservation strictly, but the energy conservation only loosely [23], our source emits several pairs of modes, as shown by the correlation peak which is elongated along the line $v_z^+ = -v_z^-$ [21].

If the pair production process is coherent, emitted pairs will be in a superposition of several pair states, each with well defined velocities. In other words, our source of atom pairs should produce pairs of entangled atoms. By filtering the velocities at the detector according to: $mv_z^+ = p$ or p' , and $mv_z^- = -p$ or $-p'$, where m is the mass of the atom, we therefore expect to obtain a Bell state of the form (1). The next step is to observe an interference between the two components of the superposition state with the interferometer in Fig. 1. This is realized using Bragg diffraction of the atoms on a second moving optical lattice oriented along the z -axis, distinct from the lattice driving the pair emission. This Bragg lattice is pulsed first for 100 μ s to realize the Bragg deflector (π -pulse), and then for 50 μ s to realize the Bragg splitters ($\pi/2$ -pulse). The reciprocal vector k_ℓ of the lattice is set to the value $\hbar k_\ell/m = 50$ mm/s. By construction, the in-




FIG. 3. Normalized cross-correlation $g^{(2)}(v_z^+, v_z^-)$. The velocities are measured along the z -axis and relative to the center-of-mass velocity of the atom pairs. A sliding average was performed to reduce the statistical noise. The correlation peak is elongated along the anti-diagonal because the source can emit in several pairs of modes. The width of the correlation peak along the diagonal corresponds to the diffraction limit imposed by the spatial extent of the source.

terferometer is closed for any pair of modes $(p, -p')$, or $(-p, p')$, satisfying the condition $p + p' = \hbar k_\ell$. The frequency difference between the laser beams forming the lattice is tuned to resonantly couple the modes with velocities $v_z^\pm = \pm 25$ mm/s but the spectral broadening induced by the finite interaction time of the atoms with the lattice is such that all pairs of modes populated in the experiment are coupled with almost the same strength. Thus, a single Bragg pulse simultaneously realizes the deflection, or the mixing, of the two pairs of modes $(p, -p')$ and $(-p, p')$.

We choose to apply the deflecting pulse right after the transfer to the $m_J = 0$ state, at $t_1 = 1100$ μ s, where the time origin is arbitrarily set at the instant when the optical lattice driving the pair emission is switched on, and t_1 is the beginning of the pulse. To close the interferometer, the time t_2 for the splitting pulse is determined experimentally. This is achieved by performing a Hong–Ou–Mandel experiment [26]; that is, we vary the time at which the splitting pulse is applied and measure the probability of joint detection at velocities $v_z^\pm = \pm 25$ mm/s (dashed lines in Fig. 1). The interferometer is closed when the joint detection probability is minimum, which, in our experiment, occurs when the Bragg splitting pulse starts at $t_2 = 1950$ μ s [21].

Ideally, one would vary the relative phase $(\phi_A - \phi_B)$ in a controlled manner to observe the modulation predicted in Eq. (10). This is not possible with the setup described here because the two splitters are realized with




FIG. 4. Joint detection probabilities measured at the output of the four-mode interferometer for three independent sets of momentum modes (p, p') . The lower graph displays the correlation coefficient, E . The gray line represents the zero level of this coefficient, calibrated using uncorrelated sets of velocities; the width of the line is the error on the zero level. The velocities v_z^\pm corresponding to the modes p are 27.0, 29.1 and 31.1 mm/s for sets 1, 2 and 3, respectively. Averages were taken over 2218 repetitions of the experiment. Error bars denote the statistical uncertainty and are obtained by bootstrapping.

the same Bragg pulse. Active control of the relative phase $(\phi_A - \phi_B)$ could be achieved using independent Bragg pulses for the splitters A and B , and we intend to implement this procedure in the future. However, we still have a way to probe different relative phases in the current setup by filtering modes for which the Bragg diffraction is slightly off-resonant, which adds a velocity-dependent contribution to the relative phase $(\phi_A - \phi_B)$ [21]. We therefore obtain different relative phases by filtering different output modes.

The number of independent sets of momentum modes $(\pm p, \pm p')$ —and thus the number of different relative phases—that we can access is constrained by the width of the Bragg resonance and the fact that the integration volume should be large enough that a significant number of atoms are detected. In the experiment we have been able to make measurements on three independent sets. We estimate the relative phases imprinted by the off-resonant Bragg diffraction to be -43° , -94° and -144° for sets 1, 2 and 3, respectively. We give these values only as an indication of how the relative phase may vary between the three sets of modes since other contributions

could also be present.

Figure 4 displays the result of these three measurements. The upper two graphs show the four joint detection probabilities [21]. As expected from Eqs. (7) and (8), the values of $\mathcal{P}(A_+, B_+)$ and $\mathcal{P}(A_-, B_-)$ on the one hand, and $\mathcal{P}(A_+, B_-)$ and $\mathcal{P}(A_-, B_+)$ on the other, appear to be correlated. Note that, for each set of modes, the sum of all four joint detection probabilities is equal to unity by construction. The lower graph shows the correlation coefficient E defined in Eq. (10). We observe that, for at least one set of modes, the correlation coefficient takes a non-zero value (set 3 gives $E = 0.51 \pm 0.20$). We can also use our data to verify the zero level of E : by filtering 78 sets of uncorrelated velocities, we find $E = -0.003$ with a statistical uncertainty of 0.020 [21] (gray line in the lower graph of Fig. 4).

Our results are thus consistent with the existence of an entangled state of two atoms. To make a stronger claim we would like to observe the modulation of E when we control the phase difference $(\phi_A - \phi_B)$. This is best achieved by introducing separate Bragg splitters. Performing a correlation measurement on a single set of momentum modes would render common any velocity dependent phase and we could then examine the variation of E with the relative phase and measure the contrast. A contrast in excess of $1/\sqrt{2}$ would permit the observation of a Bell inequality violation for freely falling massive particles using their momentum degree of freedom. Finally, we note that the setup described here can in principle be adapted to mix the mode p with p' , and $-p$ with $-p'$, by changing the reciprocal wavevector of the Bragg lattices. This variant, where the trajectories of the two atoms never cross, can also lead to a violation of a Bell inequality, in a situation where non-locality is more striking.

The research leading to these results has received funding from the European Research Council and from the People Programme (Marie Curie Actions) under the European Union’s Seventh Framework Programme (FP7/2007-2013) and H2020 Programme (2014-2020) / ERC grant agreement n°267775, and REA grant agreements n°618760 and 704832. We also acknowledge funding from the ANR through the grant agreement n°15-CE30-0017 and support from Churchill College, Cambridge.

APPENDIX

Optical lattice for pair emission

The optical lattice driving the dynamical instability at the origin of the pair creation has a period $a = 550$ nm and a depth of $0.45 E_{\text{rec}}$, where $E_{\text{rec}} = \pi^2 \hbar^2 / 2ma^2$ is the recoil energy and m is the mass of an atom. The frequency difference between the two laser beams form-

ing the lattice is $\nu = 105$ kHz, resulting in a velocity $\nu a = 57$ mm/s for the motion of the standing wave in the laboratory frame of reference.

Integration volumes for counting the atom numbers

Depending on the observable, we choose different integration volumes in velocity space in order to optimize the signal to noise ratio. In Tab. I, we summarize the integration volumes used to count the number of atoms for each graph of the main text and supplemental material.

Table I. Integration volumes. Rectangular boxes have a size δv_x , δv_y and δv_z along x , y and z , respectively. Cylindrical boxes are oriented along z ; their diameter is $\delta v_x = \delta v_y$ and their length is δv_z . All sizes are given in mm/s.

	box shape	δv_x	δv_y	δv_z
Fig. 2	rectangular	9.2	1.7	9.2
Fig. 3	cylindrical	32.0	32.0	2.5
Fig. 4	cylindrical	4.0	4.0	2.0
Fig. S1 (left)	cylindrical	18.0	18.0	1.8
Fig. S1 (right)	cylindrical	18.0	18.0	0.9
Fig. S3	cylindrical	4.0	4.0	2.6

Normalized cross-correlation

The normalized cross-correlation $g^{(2)}(v_z^+, v_z^-)$ shown in Fig. 3 of the main text displays a peak centered around $v_z^+ = -v_z^- \simeq 25$ mm/s. This peak is elongated along the line $v_z^+ = -v_z^-$, indicating the multi-mode nature of our source of atom pairs. Projections of the two-dimensional cross-correlation function along the lines $v_z^+ = -v_z^-$ and $v_z^+ - v_z^- = 50$ mm/s, corresponding to the long and short axes of the correlation peaks, respectively, are given in Fig. 5. Unlike the two-dimensional map displayed in Fig. 3 of the main text, no sliding average was performed and all experimental points are statistically independent. The different amplitudes of the correlation peak along the long and short axes stem from the different integration volumes. A Gaussian fit yields the half-widths (standard deviation) $\sigma = (9.0 \pm 2.3)$ mm/s for the long axis, and $\sigma = (2.7 \pm 0.7)$ mm/s for the short axis. These values are to be compared to the half-width of the auto-correlation functions, $\sigma_{\text{auto}} = (1.9 \pm 0.4)$ mm/s, which is the diffraction limit of our source.

Timing of the Bragg pulses

In principle, the interferometer is closed when the time t_2 at which the mixing is realized equals twice the time




FIG. 5. Projections of the two-dimensional cross-correlation function on its long and short axes. The blue points represent the experimental data. The error bars represent the statistical uncertainty and are obtained by bootstrapping. The grey lines are Gaussian fits with an offset fixed at unity.

t_1 at which the deflection is realized. However, neither the pair emission, nor the Bragg diffraction occur at a well defined instant and we have to determine experimentally the time at which the Bragg splitting pulse must be applied in order to close the interferometer. We solve this problem by performing a Hong–Ou–Mandel experiment. This is achieved by filtering two symmetric output modes, C_+ and C_- , corresponding to the input state $|p'', -p''\rangle$ (see Fig. 6). This configuration realizes a two-mode interferometer when $2p'' = \hbar k_\ell$. In the experiment we have selected the modes with the velocities $v_z^+ = -v_z^- = 25$ mm/s, which are located at the maxima of the initial velocity distribution of the emitted atom pairs. We then vary the time at which the Bragg splitting pulse is applied, and measure the probability of joint detection in the two output modes:

$$\mathcal{P}(C_+, C_-) = 2\Lambda^{-1} \langle n(p'') n(-p'') \rangle, \quad (12)$$

where the normalization factor is given by

$$\Lambda = \langle n(p'')(n(p'') - 1) \rangle + \langle n(-p'')(n(-p'') - 1) \rangle + 2\langle n(p'')n(-p'') \rangle \quad (13)$$

and we have used the notation $n(\pm p'')$ instead of $n(v_z^\pm)$, with $mv_z^\pm = \pm p''$.

In a closed interferometer, the “which-path” information is erased and the two atoms of a pair become indistinguishable after the Bragg splitter. A two-particle interference then results in the cancellation of the joint detection probability. We show the result of this measurement in Fig. 7. The dip in the joint detection probability is clearly visible when the Bragg splitting pulse is applied at time $t_2 = 1950 \mu\text{s}$, and we use this timing to realize the four-mode interferometer.




FIG. 6. Diagram of the Hong–Ou–Mandel interferometer. By filtering, only two output modes corresponding to the initial pair state $|p'', -p''\rangle$, the four-mode interferometer folds onto a two-mode interferometer. The Hong–Ou–Mandel effect occurs when the Bragg splitting pulse mixes the two input modes p'' and $-p''$. It manifests as a reduction of the probability of joint detection in the output modes C_+ and C_- , shown in Fig. 7.

Bragg diffraction model

The Bragg reflectors and splitters are realized by Bragg diffraction on a vertical, moving optical lattice formed by the interference pattern of two laser beams with slightly different frequencies. The frequency difference between the two beams forming the lattice is chosen such that the lattice is at rest in the center-of-mass reference frame of the free-falling atom pair. In the limit of a shallow lattice, i.e. when the lattice depth is smaller than the recoil energy $E_{\text{rec}} = \hbar^2 k_\ell^2 / 2m$, Bragg diffraction couples only pairs of momentum states $(p, -p')$, or $(-p, p')$, satisfying both momentum conservation: $p + p' = 2\hbar k_\ell$, and energy conservation: $p^2/2m = p'^2/2m$. If the interaction time between the atoms and the lattice is short, however, the energy conservation condition is not strict.

Resonant diffraction

We consider here the pair of input modes p and $-p'$ resonantly coupled by the Bragg lattice. We write the coupling Hamiltonian in the basis $\{|p\rangle, |-p'\rangle\}$ as:

$$\hat{H} = \frac{\hbar\Omega}{2} \begin{pmatrix} 0 & e^{i\phi} \\ e^{-i\phi} & 0 \end{pmatrix}, \quad (14)$$

where Ω is the two-photon Rabi-frequency and ϕ is the relative phase between the two laser beams forming the




FIG. 7. Joint detection probability in the two symmetric output modes as a function of the time at which the Bragg splitting pulse is applied. The blue points represent the experimental data. The error bars represent the statistical uncertainty and are obtained by bootstrapping. The grey line is a Gaussian fit. The reduction of the joint detection probability at $t_2 = 1950 \mu\text{s}$ results from the Hong–Ou–Mandel effect, and signals that the interferometer is closed.

Bragg lattice. The Bragg lattice drives a Rabi oscillation between the two modes p and $-p'$. The evolution operator describing this dynamics takes the simple form:

$$\hat{U}(t) \equiv e^{-i\hat{H}t/\hbar} = \begin{pmatrix} \cos(\Omega t/2) & -i e^{-i\phi} \sin(\Omega t/2) \\ -i e^{i\phi} \sin(\Omega t/2) & \cos(\Omega t/2) \end{pmatrix}, \quad (15)$$

where the time origin is set at the instant when the laser beams are switched on. An interaction time $t = \pi/\Omega$ (π -pulse) turns an input state $|p\rangle$ into an output state $|-p'\rangle$, and an input state $|-p'\rangle$ into an output state $|p\rangle$; it therefore realizes a Bragg deflector. Similarly, an interaction time $t = \pi/2\Omega$ ($\pi/2$ -pulse) turns $|p\rangle$ or $|-p'\rangle$ into a superposition with equal weights of $|p\rangle$ and $|-p'\rangle$; it therefore realizes a 50:50 Bragg splitter.

In our interferometer, a π -pulse and a $\pi/2$ -pulse are successively applied to realize the deflection and the splitting. Using the subscripts D and A to label the deflecting pulse and the splitting pulse A , respectively, we therefore obtain the output modes A_+ and A_- by writing:

$$\begin{pmatrix} A_+ \\ A_- \end{pmatrix} = \hat{U}_A(\pi/2\Omega) \hat{U}_D(\pi/\Omega) \begin{pmatrix} p \\ -p' \end{pmatrix} \quad (16)$$

$$= \frac{-1}{\sqrt{2}} \begin{pmatrix} e^{-i(\phi_A - \phi_D)} & i e^{-i\phi_D} \\ i e^{i\phi_D} & e^{i(\phi_A - \phi_D)} \end{pmatrix} \begin{pmatrix} p \\ -p' \end{pmatrix}. \quad (17)$$

The same reasoning applies if we consider the pair of

input modes p' and $-p$. We then obtain:

$$\begin{pmatrix} B_+ \\ B_- \end{pmatrix} = \hat{U}_B(\pi/2\Omega) \hat{U}_D(\pi/\Omega) \begin{pmatrix} p' \\ -p \end{pmatrix} \quad (18)$$

$$= \frac{-1}{\sqrt{2}} \begin{pmatrix} e^{-i(\phi_B - \phi_D)} & i e^{-i\phi_D} \\ i e^{i\phi_D} & e^{i(\phi_B - \phi_D)} \end{pmatrix} \begin{pmatrix} p' \\ -p \end{pmatrix}. \quad (19)$$

Equations (S4-S7) directly give Eqs. (2-5) in the main text.

Off-resonant diffraction

We now consider the pair of input modes p and $-p'$ for which the Bragg diffraction is slightly off-resonant. We introduce the detuning from the resonance condition: $\hbar\delta = p^2/2m - p'^2/2m$. We assume $p > p'$, so that $\delta > 0$. To first order in δ/Ω , the evolution operator in the basis $\{|p\rangle, |-p'\rangle\}$ is modified according to:

$$\hat{U}(t) \simeq \begin{pmatrix} e^{-i\delta t/2} \cos(\Omega t/2) & -i e^{-i(\phi + \delta t/2)} \sin(\Omega t/2) \\ -i e^{i(\phi + \delta t/2)} \sin(\Omega t/2) & e^{i\delta t/2} \cos(\Omega t/2) \end{pmatrix}. \quad (20)$$

If we consider instead the input states p' and $-p$, but keep the same definition for δ , we must take care to replace δ by $-\delta$ in this evolution operator. Compared to the resonant case, one sees that an additional phase δt is accumulated between the components $|p\rangle$ and $|-p'\rangle$ during the interaction with the Bragg lattice. At the output of the interferometer, the modes A_\pm and B_\pm are now given by the matrix equations

$$\begin{pmatrix} A_+ \\ A_- \end{pmatrix} \simeq \frac{-1}{\sqrt{2}} \begin{pmatrix} e^{-i(\phi_A - \phi_D - \pi\delta/4\Omega)} & i e^{-i(\phi_D + 3\pi\delta/4\Omega)} \\ i e^{i(\phi_D + 3\pi\delta/4\Omega)} & e^{i(\phi_A - \phi_D - \pi\delta/4\Omega)} \end{pmatrix} \begin{pmatrix} p \\ -p' \end{pmatrix} \quad (21)$$

and

$$\begin{pmatrix} B_+ \\ B_- \end{pmatrix} \simeq \frac{-1}{\sqrt{2}} \begin{pmatrix} e^{-i(\phi_B - \phi_D + \pi\delta/4\Omega)} & i e^{-i(\phi_D - 3\pi\delta/4\Omega)} \\ i e^{i(\phi_D - 3\pi\delta/4\Omega)} & e^{i(\phi_B - \phi_D + \pi\delta/4\Omega)} \end{pmatrix} \begin{pmatrix} p' \\ -p \end{pmatrix}. \quad (22)$$

Inverting the matrix equations (21) and (22), we can express the entangled state $|\psi\rangle = \frac{1}{\sqrt{2}}(|p, -p\rangle + |p', -p'\rangle)$ at the output of the interferometer:

$$|\Psi_{\text{out}}\rangle \simeq \frac{1}{2\sqrt{2}} \left[-i(e^{i(\phi_A - \pi\delta/\Omega)} + e^{i(\phi_B + \pi\delta/\Omega)})|A_+, B_+\rangle \right. \\ \left. + (e^{i(\phi_A - \phi_B - \pi\delta/2\Omega)} - e^{3i\pi\delta/2\Omega})|A_+, B_-\rangle \right. \\ \left. + (e^{-i(\phi_A - \phi_B - \pi\delta/2\Omega)} - e^{-3i\pi\delta/2\Omega})|A_-, B_+\rangle \right. \\ \left. - i(e^{-i(\phi_A - \pi\delta/\Omega)} + e^{-i(\phi_B + \pi\delta/\Omega)})|A_-, B_-\rangle \right]. \quad (23)$$

We finally obtain the joint detection probabilities

$$\mathcal{P}(A_\pm, B_\pm) \simeq \frac{1}{2} \cos^2 [(\phi_A - \phi_B - 2\pi\delta/\Omega)/2], \quad (24)$$

$$\mathcal{P}(A_\pm, B_\mp) \simeq \frac{1}{2} \sin^2 [(\phi_A - \phi_B - 2\pi\delta/\Omega)/2]. \quad (25)$$

To first order in δ/Ω , the effect of the detuning from the Bragg resonance is thus to add a contribution $-2\pi\delta/\Omega$ to the relative phase $(\phi_A - \phi_B)$. This off-resonance contribution depends on p and p' through the detuning δ .

The velocities v_z^\pm corresponding to the mode p are 27.0, 29.1 and 31.1 mm/s for sets 1, 2 and 3, respectively. The velocities corresponding to the mode p' are 23.0, 20.9 and 18.9 mm/s for sets 1, 2 and 3, respectively. The detuning from the Bragg resonance condition for these three sets of modes are thus: $\delta_1/2\pi = 0.9$ kHz, $\delta_2/2\pi = 1.9$ kHz and $\delta_3/2\pi = 2.9$ kHz. For these values of the detuning δ , the condition $\delta \ll \Omega$ is only marginally satisfied and the lowest order approximation overestimates the relative phases by about 30%. For a better estimation, we wrote the exact evolution operator for the two-mode dynamics, and numerically calculated the additional phases with respect to the resonant case; we found -43 , -94 and -144° for sets 1, 2 and 3, respectively.

Experimental measurement of the joint detection probabilities

The probabilities of joint detection in the output modes A_\pm and B_\pm are measured by counting the number of atoms with velocities $mv_z^+ = p$ or p' , and $mv_z^- = -p$ or $-p'$, and using the relations

$$\mathcal{P}(A_+, B_+) = \Lambda^{-1} \langle n(p) n(p') \rangle, \quad (26)$$

$$\mathcal{P}(A_-, B_-) = \Lambda^{-1} \langle n(-p) n(-p') \rangle, \quad (27)$$

$$\mathcal{P}(A_+, B_-) = \Lambda^{-1} \langle n(p) n(-p) \rangle, \quad (28)$$

$$\mathcal{P}(A_-, B_+) = \Lambda^{-1} \langle n(-p') n(p') \rangle, \quad (29)$$

where the normalization factor is given by

$$\Lambda = \langle n(p) n(p') \rangle + \langle n(-p) n(-p') \rangle \\ + \langle n(p) n(-p) \rangle + \langle n(-p') n(p') \rangle \quad (30)$$

and we have used the notation $n(\pm p)$ or $n(\pm p')$ instead of $n(v_z^\pm)$, with $mv_z^\pm = \pm p$ or $\pm p'$.

Zero level of the correlation coefficient

For measuring the joint detection probabilities, we have counted the number of atoms $n(v_z^\pm)$ detected at four different velocities: $mv_z^+ = p$ or p' and $mv_z^- = -p$ or $-p'$. This was repeated with three different sets of modes $(\pm p, \pm p')$ for which the interferometer was closed.

In order to confirm the zero level of the correlation coefficient E , we have constructed a correlation coefficient using sets of modes which are initially uncorrelated, and do not close the interferometer. Using the $3^4 = 81$ values of v_z^\pm , there are $81 - 3 = 78$ such combinations. The mean values of the joint detection probabilities measured with these uncorrelated sets of modes are all close to $1/4$,




FIG. 8. Histogram of the correlation coefficient measured with 78 sets of uncorrelated modes, which do not close the interferometer. The fact that the distribution is peaked around zero shows that there is no bias in the evaluation of the correlation coefficient.

as summarized in Tab. II. In Fig. 8, we show a histogram of the corresponding values of E . The distribution has a mean value of -0.003 ± 0.020 . These calibration measurements give us confidence that we have no systematic bias in the estimation of the correlation coefficient.

Table II. Mean values of the joint detection probabilities for uncorrelated data.

$\mathcal{P}(A_+, B_+)$	0.245 ± 0.007
$\mathcal{P}(A_-, B_-)$	0.257 ± 0.010
$\mathcal{P}(A_+, B_-)$	0.258 ± 0.009
$\mathcal{P}(A_-, B_+)$	0.241 ± 0.009

- [1] J. P. Dowling and G. J. Milburn, *Phil. Trans. R. Soc. Lond. A* **361**, 1655 (2003).
- [2] A. Aspect, “Introduction: John bell and the second quantum revolution,” in *Speakable and Unsayable in Quantum Mechanics: Collected Papers on Quantum Philosophy* (Cambridge University Press, Cambridge, 2004) pp. xvii–xl.
- [3] R. P. Feynman, *Int. J. Theor. Phys.* **21**, 467 (1982).
- [4] N. Gisin and M. Rigo, *J. Phys. A: Math. Gen.* **28**, 7375 (1995).
- [5] A. Aspect, “Bell’s Theorem: The Naive View of an Experimentalist,” in *Quantum (Un)speakables: From Bell*

- to Quantum Information* (Springer, Berlin, Heidelberg, 2002) pp. 119–153.
- [6] M. C. Tichy, F. Mintert, and A. Buchleitner, *J. Phys. B: At. Mol. Opt.* **44**, 192001 (2011).
- [7] J. S. Bell, *Physics* **1**, 195 (1964).
- [8] A. Einstein, B. Podolsky, and N. Rosen, *Phys. Rev.* **47**, 777 (1935).
- [9] U. Fano, *Am. J. Phys.* **29**, 539 (1961).
- [10] R. J. Glauber, in *Quantum Optics and Electronics*, edited by C. de Witt, A. Blandin, and C. Cohen-Tannoudji (Gordon and Breach, New York, 1965).
- [11] C. K. Hong, Z. Y. Ou, and L. Mandel, *Phys. Rev. Lett.* **59**, 2044 (1987).
- [12] The requirement for four modes holds for systems of two particles. In the context of continuous variables, configurations involving only two modes can also lead to violations of Bell’s inequalities (see, for instance, [27, 28]).
- [13] A. Aspect, *Nature* **398**, 189 (1999).
- [14] A. Aspect, *Physics* **8**, 123 (2015).
- [15] A two-electron interference in four momentum modes was reported in Ref. [29], but without access to all four joint detection probabilities.
- [16] R. Penrose, *Gen. Relat. Gravit.* **28**, 581 (1996).
- [17] M. A. Horne, A. Shimony, and A. Zeilinger, *Phys. Rev. Lett.* **62**, 2209 (1989).
- [18] J. G. Rarity and P. R. Tapster, *Phys. Rev. Lett.* **64**, 2495 (1990).
- [19] J. Kofler, M. Singh, M. Ebner, M. Keller, M. Kotyrbá, and A. Zeilinger, *Phys. Rev. A* **86**, 032115 (2012).
- [20] R. J. Lewis-Swan and K. V. Kheruntsyan, *Phys. Rev. A* **91**, 052114 (2015).
- [21] See Appendix for a model of Bragg diffraction and details of other methods.
- [22] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Phys. Rev. Lett.* **23**, 880 (1969).
- [23] M. Bonneau, J. Ruaudel, R. Lopes, J.-C. Jaskula, A. Aspect, D. Boiron, and C. I. Westbrook, *Phys. Rev. A* **87**, 061603 (2013).
- [24] This number refers to the average atom number in the integration volumes used in Fig. 4. It takes into account the estimated 25% quantum efficiency of the detector.
- [25] M. Schellekens, R. Hoppeler, A. Perrin, J. V. Gomes, D. Boiron, A. Aspect, and C. I. Westbrook, *Science* **310**, 648 (2005).
- [26] R. Lopes, A. Imanaliev, A. Aspect, M. Cheneau, D. Boiron, and C. I. Westbrook, *Nature* **520**, 66 (2015).
- [27] J. Wenger, M. Hafezi, F. Grosshans, R. Tualle-Brouiri, and P. Grangier, *Phys. Rev. A* **67**, 012105 (2003).
- [28] D. Cavalcanti, N. Brunner, P. Skrzypczyk, A. Salles, and V. Scarani, *Phys. Rev. A* **84**, 022105 (2011).
- [29] M. Waitz, D. Metz, J. Lower, C. Schober, M. Keiling, M. Pitzer, K. Mertens, M. Martins, J. Viefhaus, S. Klumpp, T. Weber, H. Schmidt-Böcking, L. Schmidt, F. Morales, S. Miyabe, T. Rescigno, C. McCurdy, F. Martín, J. Williams, M. Schöffler, T. Jahnke, and R. Dörner, *Phys. Rev. Lett.* **117**, 083002 (2016).